

# Mechanisms of Extreme Precipitation Change Over the Northeast

**Jonathan Winter<sup>1,2</sup>, Huanping Huang<sup>2,1</sup>,  
Erich Osterberg<sup>2</sup>**

<sup>1</sup>Department of Geography, Dartmouth College,



<sup>2</sup>Department of Earth Sciences, Dartmouth College




**BREE**  
Basin Resilience to  
Extreme Events  
in the Lake Champlain Basin




# Northeast Extreme Precipitation Events Increased Dramatically 1960-Present




# Northeast Extreme Precipitation Increased Dramatically ~~1960-1996~~ 1996-Present



# Northeast Extreme Precipitation Increase is Widespread




# Blame Tropical Cyclones (and Fronts and Extratropical Cyclones)





# GHCN-D, LI2013, NARR over the Northeast

- Global Historical Climatology Network – Daily: GHCN-D; Menne et al. 2012; 1901-2014
- Gridded observations of Livneh et al.: LI2013; Livneh et al. 2013; 1915-2011
- North American Regional Reanalysis: NARR; Mesinger et al. 2006; 1979-2014





# Trends Sensitive to Start Year, Changepoint Analysis Better Characterizes Time Series

- Trends generally increasing with later start year: 2.4 mm decade<sup>-1</sup> (1901-2014) to 14.7 mm decade<sup>-1</sup> (1979-2014)
- Change in extreme best characterized as a shift in 1996: 53%




# Extreme Precipitation: Averages Higher on Coasts, Trends Consistent over Domain

- Annual extreme precipitation ( $\text{mm decade}^{-1}$ ) increased in 58 (50%), 30 (25%) of which were statistically significant
- Five (4.3%) stations had negative trends, two of which were significant






# Extreme Precipitation: Difference between 1996-2014 and 1901-1995

- Annual extreme precipitation (%) was higher in 105 stations (91%) after 1996, with 56 stations exceeding a 50% increase
- Decreases east of Lake Erie (western NY and PA) and northeast WV




# Extreme Precipitation Trends Most Difficult for Gridded Observations and Reanalysis

- LI2013 reproduces GHCN-D extreme precipitation, but underestimates GHCN-D extreme precipitation trends
- NARR underestimates GHCN-D extreme precipitation, and NARR extreme precipitation trends are sensitive to end year



# Why Has Extreme Precipitation Increased?


- Global Historical Climatology Network – Daily: GHCN-D; Menne et al. 2012; 1979-2016
- US Daily Weather Maps: NOAA 2017; 1979-2016
- ERA Interim Reanalysis: ERA-I; Dee et al. 2011, 1979-2016




# Focus on Early Fall (Sep-Oct), Early Summer (Jun-Jul), and Late Winter (Feb-Mar)

Month	Pre-1996 EP mean (mm yr <sup>-1</sup> )	Post-1996 EP mean (mm yr <sup>-1</sup> )	Absolute EP change (mm yr <sup>-1</sup> )	Relative EP change (%)
1	1.7	2.2	0.5	29
2	1.7	2.2	0.5	29
3	3.2	4.9	1.7	53
4	3.6	4.6	1	28
5	4.6	5.7	1.1 <sup>#</sup>	24
6	7.8	10.9	3.1	40
7	9.8	14.3	4.5 <sup>*</sup>	46
8	14.7	12.6	-2.1 <sup>#</sup>	-14
9	9.9	20.3	10.4 <sup>*</sup>	105
10	6.8	12.3	5.5	81
11	6.1	6.4	0.3	5
12	3.2	4.1	0.9	28
Total	73.1	100.5	27.4 <sup>*</sup>	37

# Categorize 273 5+ Station Extreme Precipitation Events (by Huanping)




# Sep-Oct Tropical Cyclones Dominate Northeast Extreme Precipitation Increase




# Analyze Atmospheric Drivers of the Extreme Precipitation Increase

- Composite potential driving fields on extreme precipitation days for pre-1996 (1979-1995) and post-1996 (1996-2016)
  - Total column water vapor
  - Geopotential height at 500 hPa
  - Sea level pressure
  - Zonal and meridional winds at 850 hPa and 250 hPa
  - Vertical velocity
  - Sea surface temperature (SST)




# Tropical Cyclone Extreme Precipitation Correlated with Sea Surface Temperatures


SST: Post- & Pre-1996 All Difference



# Why Less Extreme Precipitation During Last AMO+ Phase (1928-1962)?



# Increased Total Column Water Vapor in Hurricane Main Development Region



# Conclusions

- Recent increases in Northeast extreme precipitation are best characterized as an abrupt shift of 53% after 1996
- Extreme precipitation increases are distributed uniformly throughout the Northeast except western NY/PA and WV
- There are differences between GHCN-D and NARR, LI2013 extreme precipitation trends
- Eighty-eight percent of the abrupt 1996 extreme precipitation increase is explained by 5+ station events in early fall, early summer, and late winter
- The 1996 extreme precipitation increase is caused by tropical cyclones (48%), fronts (25%), and extratropical cyclones (15%)
- Increased extreme precipitation is associated with warmer Atlantic sea surface temperatures and more water vapor